



Anti-CXCL12 polyclonal antibody (CPBT-68002GM)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview Goat anti Mouse SDF-1 alpha antibody recognizes mouse SDF-1 alpha, otherwise known as CXCL12a, a stromal cell derived CXC chemokine which, together with SDF-1 beta, arises from alternative splicing of the SDF-1 gene, and acts as a chemoattractant for T and B lymphocytes, monocytes and migratory neurons. SDF-1 binds with high-affinity to the G protein-coupled receptor CXCR4 (fusin), and acts as an inhibitor of the CXCR4-mediated entry of HIV-1 virus into target T-cells. ELISA This antibody may be used in an indirect ELISA or as a detection reagent in a sandwich ELISA with AAM64 as the capture antibody and PMP77 as the standard.

| | |
|---------------------------|---|
| Specificity | SDF-1 ALPHA |
| Isotype | IgG |
| Source/Host | Goat |
| Species Reactivity | Mouse |
| Conjugate | Biotin |
| Applications | ELISA; WB |
| Format | Purified IgG conjugated to Biotin - lyophilised |
| Size | 50 µg |
| Preservative | None |
| Storage | in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |

GENE INFORMATION

| | |
|----------------------------|---|
| Gene Name | Cxcl12 chemokine (C-X-C motif) ligand 12 [Mus musculus (house mouse)] |
| Official Symbol | CXCL12 |
| Synonyms | CXCL12; chemokine (C-X-C motif) ligand 12; Pbsf; Sdf1; Tlsf; Tpar1; Scyb12; stromal cell-derived factor 1; pre-B-cell growth-stimulating factor; thymic lymphoma cell-stimulating factor; 12-O-tetradecanoylphorbol 13-acetate repressed protein 1; SDF-1 ALPHA |
| Entrez Gene ID | 20315 |
| Protein Refseq | NP_001012495 |
| UniProt ID | P40224 |
| Chromosome Location | 6 F1; 6 54.81 cM |
| Pathway | Axon guidance; Chemokine receptors bind chemokines; Chemokine signaling pathway; Class A/1 (Rhodopsin-like receptors); Cytokine-cytokine receptor interaction; Defective ACTH causes Obesity and Pro-opiomelanocortinin deficiency (POMCD); Disease; G alpha (i) signalling events; |
| Function | CXCR chemokine receptor binding; chemokine activity; chemokine receptor binding; cytokine activity; growth factor activity; |