



User's Manual

Influenza A H1N1 (Swine Flu 2009) Hemagglutinin Antibody Pair Set

REF

ABPR-ZB001



5 Plates, 15 Plates

RUO

This product is for research use only and is not intended for diagnostic use.

For illustrative purposes only. To perform the assay the instructions for use provided with the kit have to be used.

Creative Diagnostics

 Address: 45-1 Ramsey Road, Shirley, NY 11967, USA

 Tel: 1-631-624-4882 (USA) 44-161-818-6441 (Europe)  Fax: 1-631-938-8221

 Email: info@creative-diagnostics.com  Web: www.creative-diagnostics.com

PRODUCT INFORMATION

Intended Use

Quantitative determination of Influenza Hemagglutinin

General Description

ABPR-ZB001 is a solid phase sandwich ELISA for quantitative determination of Influenza Hemagglutinin.

Reagents And Materials Provided

1. Capture Antibody: 1 mg/mL of mouse anti-Influenza A H1N1 (Swine Flu 2009) Hemagglutinin monoclonal antibody [CABT-ZB428] (in PBS, pH 7.4).
2. Detection Antibody: 0.2 mg/mL of rabbit anti-Influenza A H1N1 (Swine Flu 2009) Hemagglutinin polyclonal antibody conjugated to HRP (in PBS, 50 % HRP-Protector, pH 7.4, store at 4°C).
3. Standard: Each vial contains 67 ng of recombinant Influenza A H1N1 (Swine Flu 2009) Hemagglutinin.

Reconstitution And Storage

Reconstitution

1. Capture Antibody: Dilute to a working concentration of 2 µg/mL in PBS before coating.
2. Detection Antibody: Dilute to working concentration of 1 µg/mL in detection antibody dilution buffer before use.
3. Standard: Reconstitute with 1 mL detection antibody dilution buffer.

Storage

1. Capture Antibody: Aliquot and store at -20°C to -80°C for up to 6 months from date of receipt. Avoid repeated freeze-thaw cycles.
2. Detection Antibody: Store at 4°C and protect it from prolonged exposure to light for up to 6 months from date of receipt. DO NOT FREEZE!
3. Standard: Store lyophilized standard at -20°C to -80°C for up to 6 months from date of receipt. Aliquot and store the reconstituted Standard at -80°C for up to 1 month. Avoid repeated freeze-thaw cycles.

Detection Range

Assay range: 39.06-2500 pg/mL