



# Anti-TP53 monoclonal antibody, clone DO-13 (CABT-50215MH)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Mouse anti Human p53 (aa26-35) antibody, clone DO-13 recognizes the human p53 tumor suppressor protein, binding to both wild type and mutant forms. Mouse anti Human p53 (aa26-35) antibody, clone DO-13 recognizes an epitope at the N-terminal end of p53 between amino acids 26-35.
<b>Specificity</b>	TP53
<b>Isotype</b>	IgG1
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	DO-13
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	IHC-Fr; IHC-P
<b>Format</b>	Purified IgG - liquid
<b>Size</b>	20 µg
<b>Preservative</b>	0.09% Sodium Azide
<b>Storage</b>	in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">TP53 tumor protein p53 [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	TP53
<b>Synonyms</b>	TP53; tumor protein p53; P53; BCC7; LFS1; TRP53; cellular tumor antigen p53; antigen NY-CO-13; tumor protein 53; phosphoprotein p53; p53 tumor suppressor; mutant tumor protein 53; transformation-related protein 53;
<b>Entrez Gene ID</b>	<a href="#">7157</a>
<b>Protein Refseq</b>	<a href="#">NP_000537</a>
<b>UniProt ID</b>	P04637
<b>Chromosome Location</b>	17p13.1
<b>Pathway</b>	AMPK signaling; Activation of BH3-only proteins; Activation of NOXA and translocation to mitochondria; Activation of PUMA and translocation to mitochondria; Alzheimers Disease; Amyotrophic lateral sclerosis (ALS); Apoptosis; Aurora A signaling;
<b>Function</b>	ATP binding; DNA binding; MDM2/MDM4 family protein binding; RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; RNA polymerase II core promoter sequence-specific DNA binding; RNA polymerase II transcription factor binding; RNA polymerase II transcription regulatory region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; chaperone binding; chromatin binding; copper ion binding; core promoter sequence-specific DNA binding; damaged DNA binding; enzyme binding; histone acetyltransferase binding; histone deacetylase regulator activity; identical protein binding; p53 binding; protease binding; protein N-terminus binding; protein binding; protein heterodimerization activity; protein kinase binding; protein phosphatase 2A binding; protein phosphatase binding; receptor tyrosine kinase binding; sequence-specific DNA binding RNA polymerase II transcription factor activity; sequence-specific DNA binding transcription factor activity; transcription factor binding; transcription regulatory region DNA binding; ubiquitin protein ligase binding; zinc ion binding;