



Anti-TP53 monoclonal antibody, clone FP3-2 (CABT-50221MH)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Mouse anti Human p53 (pSer392) antibody, clone FP3-2 recognizes human p53, binding only to p53 when phosphorylated at the Ser392 residue. Ser392 phosphorylation of p53 has been shown to occur following DNA damage. Phosphorylation of p53 at this site is mediated by the casein kinase 2.
Specificity	TP53
Immunogen	Synthetic peptide RKKLMFKTEGPDS[P]D.
Isotype	lgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	FP3-2
Conjugate	Unconjugated
Applications	IP; IHC-P; WB
Format	Purified IgG - liquid
Size	100 μg
Preservative	0.09% Sodium Azide
Storage	in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

GENE INFORMATION

Gene Name	TP53 tumor protein p53 [Homo sapiens (human)]
Official Symbol	TP53
Synonyms	TP53; tumor protein p53; P53; BCC7; LFS1; TRP53; cellular tumor antigen p53; antigen NY-CO-13; tumor protein 53; phosphoprotein p53; p53 tumor suppressor; mutant tumor protein 53; transformation-related protein 53;
Entrez Gene ID	<u>7157</u>
Protein Refseq	NP 000537
UniProt ID	P04637
Chromosome Location	17p13.1
Pathway	AMPK signaling; Activation of BH3-only proteins; Activation of NOXA and translocation to mitochondria; Activation of PUMA and translocation to mitochondria; Alzheimers Disease; Amyotrophic lateral sclerosis (ALS); Apoptosis; Aurora A signaling;
Function	ATP binding; DNA binding; MDM2/MDM4 family protein binding; RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; RNA polymerase II core promoter sequence-specific DNA binding; RNA polymerase II transcription factor binding; RNA polymerase II transcription regulatory region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; chaperone binding; chromatin binding; copper ion binding; core promoter sequence-specific DNA binding; damaged DNA binding; enzyme binding; histone acetyltransferase binding; histone deacetylase regulator activity; identical protein binding; p53 binding; protease binding; protein N-terminus binding; protein binding; protein heterodimerization activity; protein kinase binding; protein phosphatase 2A binding; protein phosphatase binding; receptor tyrosine kinase binding; sequence-specific DNA binding RNA polymerase II transcription factor activity; sequence-specific DNA binding transcription factor activity; transcription factor binding; transcription regulatory region DNA binding; ubiquitin protein ligase binding; zinc ion binding;