



Anti-HCAR2 (aa 250-350) polyclonal antibody (CABT-BL1761)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Synthetic peptide conjugated to KLH derived from within residues 250 - 350 of Human Puma gamma.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, ICC/IF
Cellular Localization	Cell Membrane
Format	Liquid
Size	100 µg
Buffer	1% BSA, PBS, pH 7.4
Preservative	0.02% Sodium Azide
Storage	Store at +4°C short term (1-2 weeks). Aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

BACKGROUND

Introduction

Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid> acifran> 5-methyl nicotinic acid = acipimox>> nicotinuric acid = nicotinamide.

GENE INFORMATION

Entrez Gene ID	338442
----------------	------------------------

Protein Refseq	NP_808219
----------------	---------------------------

UniProt ID	Q8TDS4
------------	------------------------
