



Anti-HIST1H2AL (aa 100-130) polyclonal antibody (CABT-BL1809)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Synthetic peptide conjugated to KLH, corresponding to a region within C terminal amino acids 100-130 of Human HIST1H2AL.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, IHC-P
Cellular Localization	Nucleus. Chromosome.
Format	Liquid
Buffer	PBS
Preservative	0.09% Sodium Azide
Storage	Store at 4°C (up to 6 months). For long term storage store at -20°C

BACKGROUND

Introduction Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B,

H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

GENE INFORMATION

Entrez Gene ID	8332
Protein Refseq	<u>NP_003502</u>
UniProt ID	A4FTV9