



Anti-MLLT4 (N-terminal) polyclonal antibody (CABT-BL2454)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Synthetic peptide: CKLPGDDRLMKNRADHRS conjugated to maleimide-activated BSA, corresponding to amino acids 1163-1179 of Rat I + S Afadin (with N-terminal added cysteine).
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Mouse, Rat, Dog, Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, IF
Cellular Localization	Cell Membrane and Nuclear
Format	Liquid
Buffer	1% BSA, 0.01M PBS, pH 7.4
Preservative	15mM Sodium Azide
Storage	Store at +4°C short term (1-2 weeks). Aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

BACKGROUND

Introduction This gene encodes a transcriptional coactivator that plays an essential role in regulating gene

expression during early development and hematopoiesis. The encoded protein contains

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

multiple conserved functional domains. One of these domains, the SET domain, is responsible for its histone H3 lysine 4 (H3K4) methyltransferase activity which mediates chromatin modifications associated with epigenetic transcriptional activation. This protein is processed by the enzyme Taspase 1 into two fragments, MLL-C and MLL-N. These fragments reassociate and further assemble into different multiprotein complexes that regulate the transcription of specific target genes, including many of the HOX genes. Multiple chromosomal translocations involving this gene are the cause of certain acute lymphoid leukemias and acute myeloid leukemias. Alternate splicing results in multiple transcript variants.

GENE INFORMATION

Entrez Gene ID	4301
Protein Refseq	NP_001035089
UniProt ID	<u>P55196</u>