



## Anti-MMP1 (full length) polyclonal antibody (CABT-BL2463)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

| Immunogen             | Full length Human MMP1  |
|-----------------------|---|
| Isotype               | IgG   |
| Source/Host           | Mouse   |
| Species Reactivity    | Human   |
| Purification          | Protein A purified  |
| Conjugate             | Unconjugated  |
| Applications          | WB  |
| Cellular Localization | Secreted; extracellular space; extracellular matrix.  |
| Format                | Liquid  |
| Buffer                | 1X PBS, pH 7.2  |
| Preservative          | None  |
| Storage               | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |

## **BACKGROUND**

**Introduction** Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of

extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

metastasis. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes a secreted enzyme which breaks down the interstitial collagens, types I, II, and III. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Mar 2009]

## **GENE INFORMATION**

| Entrez Gene ID | 4312          |
|----------------|---------------|
| Protein Refseq | NP_001139410  |
| UniProt ID     | <u>B4DN15</u> |