



Anti-NCF4 (C-terminal) polyclonal antibody (CABT-BL2583)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Synthetic peptide selected from the C terminal region of Human NCF4 conjugated to KLH.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Ammonium Sulphate Precipitation
Conjugate	Unconjugated
Applications	WB, ELISA
Cellular Localization	Cytoplasm.
Format	Liquid
Buffer	PBS
Preservative	0.09% Sodium Azide
Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

BACKGROUND

Introduction

The protein encoded by this gene is a cytosolic regulatory component of the superoxide-producing phagocyte NADPH-oxidase, a multicomponent enzyme system important for host defense. This protein is preferentially expressed in cells of myeloid lineage. It interacts primarily with neutrophil cytosolic factor 2 (NCF2/p67-phox) to form a complex with neutrophil cytosolic

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

factor 1 (NCF1/p47-phox), which further interacts with the small G protein RAC1 and translocates to the membrane upon cell stimulation. This complex then activates flavocytochrome b, the membrane-integrated catalytic core of the enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests its role in PI(3) kinase-mediated signaling events. The phosphorylation of this protein was found to negatively regulate the enzyme activity. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]

GENE INFORMATION

Entrez Gene ID	<u>4689</u>
Protein Refseq	NP_000622
UniProt ID	<u>Q15080</u>