



Anti-OR2M2 (internal region) polyclonal antibody (CABT-BL2804)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Synthetic peptide derived from an internal sequence within human OR2M2.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, ELISA, ICC/IF
Cellular Localization	Cell membrane.
Format	Liquid
Buffer	50% Glycerol, PBS (without Mg2+ and Ca2+), 150mM Sodium chloride, pH 7.4
Preservative	0.02% Sodium Azide
Storage	Store at -20°C. Stable for 12 months at -20°C

BACKGROUND

Introduction

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

GENE INFORMATION

Entrez Gene ID	<u>391194</u>
Protein Refseq	NP_001004688
UniProt ID	Q96R28