



Anti-TXNDC3 (full length) polyclonal antibody (CABT-BL3711)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Full length TXNDC3 protein (Human)
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Purification	Protein A purified
Conjugate	Unconjugated
Applications	WB
Cellular Localization	Cytoplasm
Format	Liquid
Buffer	PBS, pH 7.2
Preservative	None
Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

BACKGROUND

Introduction This gene encodes a protein with an N-terminal thioredoxin domain and three C-terminal

nucleoside diphosphate kinase (NDK) domains, but the NDK domains are thought to be

catalytically inactive. The sea urchin ortholog of this gene encodes a component of sperm outer

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

dynein arms, and the protein is implicated in ciliary function. Mutations in this gene are implicated in primary ciliary dyskinesia type 6.[provided by RefSeq, Nov 2009]

GENE INFORMATION

Entrez Gene ID	51314
Protein Refseq	NP_057700
UniProt ID	Q8N427