



Chicken Anti-Avian ADV Group II (HEV) Polyclonal antibody (CABT-CS207)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| Specificity | ADV |
|--------------------|-----------------|
| Target | ADV |
| Isotype | IgY |
| Source/Host | Chicken |
| Species Reactivity | ADV |
| Conjugate | unconjugated |
| Format | Liquid |
| Size | 1 ml |
| Buffer | Antiserum |
| Preservative | None |
| Storage | Store at -20°C. |

BACKGROUND

Introduction

The avian adenoviruses can be divided into three groups, i.e. groups I, II and III. Group I, or conventional adenoviruses, share a common group antigen, distinct from the mammalian adenovirus group antigen. Group II adenoviruses include the viruses of turkey haemorrhagic enteritis (THE), marble spleen disease (MSD) and group II splenomegaly of chickens. These viruses share a common antigen which is distinct from the group antigen of mammalian and

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

group I avian adenoviruses. Group III viruses, the egg drop syndrome (EDS) viruses, are widely distributed in waterfowl but can easily infect chickens, resulting in the production of abnormal eggshells.

Keywords

Avian Adenovirus; Avian ADV; ADV; Adenovirus; Hemorrhagic enteritis virus; HEV