



Mouse anti Canine IL10 monoclonal antibody, clone 249222 (CABT-L123)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| | |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Specificity | Delects canine IL-10 in ELISAs. In ELISAs, this antibody does not cross-react with recombinant feline IL-10, recombinant human (rh) IL-10, recombinant mouse (rm) IL-10, recombinant porcine IL-10, rhIL-10 sR, rmIL-10 sR, or recombinant rat IL-10. |
| Target | IL-10 |
| Immunogen | E. coli-derived recombinant canine IL-10, Ser20-Ile179, Accession #XP_855560 |
| Isotype | IgG1 |
| Source/Host | Mouse |
| Species Reactivity | Canine |
| Clone | 249222 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Conjugate | Unconjugated |
| Applications | ELISA(Cap) |
| Reconstitution | Reconstitute at 0.5 mg/mL in sterile PBS. |
| Format | Lyophilized; Small package size(SP): Liquid |
| Size | 25 µg, 500 µg |
| Buffer | PBS with Trehalose |
| Preservative | None |

| | |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution. |
| Ship | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C. |

BACKGROUND

| | |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Introduction | IL-10, initially designated cytokine synthesis inhibitory factor (CSIF), was originally identified as a product of murine T helper 2 (Th2) clones that inhibited the cytokine production by Th1 clones. The canine homologue of mouse IL-10 has been cloned. Canine IL-10 shares 80% amino acid sequence homology with human IL-10 and 72% sequence homology with mouse IL-10. IL-10 is a pleiotropic cytokine that can exert either immunostimulatory or immunosuppressive effects on a variety of cell types. It is a potent immunosuppressant of macrophage functions. In vitro, IL-10 can inhibit the accessory function and antigen-presenting capacity of monocytes by, among other effects, downregulating class II MHC expression. Thus, IL-10 can inhibit monocyte/macrophage-dependent, antigen stimulated cytokine synthesis (especially IFN-gamma) by human PBMNC and NK, and mouse Th1 cells. Additionally, IL-10 is a potent inhibitor of monocyte/macrophage activation and its resultant cytotoxic effects. As an immunostimulatory cytokine, IL-10 can act on B cells to enhance their viability, cell proliferation, Ig secretion, and class II MHC expression. Aside from B lymphocytes, IL-10 is also a growth co-stimulator for thymocytes and mast cells, as well as an enhancer of cytotoxic T cell development. |
| Keywords | CSIF;CSIFMGC126450;Cytokine synthesis inhibitory factor;IL10;IL-10;IL10A;IL-10MGC126451;interleukin 10;interleukin-10;TGIF |

GENE INFORMATION

| | |
|-----------------------|------------------------|
| Entrez Gene ID | 403628 |
| UniProt ID | P48411 |