



Goat anti Human IL1B polyclonal antibody [Biotin] (CABT-L199)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	Detects human IL-1 beta /IL-1F2 in ELISAs and Western blots. In sandwich immunoassays, less than 4% cross-reactivity with recombinant ratIL-1 beta is observed and less than 0.1% cross-reactivity with recombinant mouse IL-1 beta and recombinant porcine IL-1 beta is observed.
Target	IL-1beta/IL-1F2
Immunogen	E. coli-derived recombinant human IL-1 beta /IL-1F2, Ala117-Ser269, Accession #NP_000567
Isotype	IgG
Source/Host	Goat
Species Reactivity	Human
Purification	Antigen Affinity-purified
Conjugate	Biotin
Applications	ELISA(Det), ICC/IF, WB
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Format	Lyophilized
Size	50 µg
Buffer	PBS with BSA
Preservative	None

Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.
Ship	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

BACKGROUND

Introduction	IL-1 is a name that designates two pleiotropic cytokines, IL-1 alpha(IL-1F1) and IL-1 beta(IL-1F2), which are the products of distinct genes. IL-1 alpha and IL-1 beta are structurally related polypeptides that share approximately 21% amino acid (aa) identity in human. Both proteins are produced by a wide variety of cells in response to inflammatory agents, infections, or microbial endotoxins. While IL-1 alpha and IL-1 beta are regulated independently, they bind to the same receptor and exert identical biological effects. IL-1 RI binds directly to IL-1 alpha or IL-1 beta b and then associates with IL-1 R accessory protein (IL-1 R3/IL-1 R AcP) to form a high-affinity receptor complex that is competent for signal transduction. IL-1 RII has high affinity for IL-1 beta but functions as a decoy receptor and negative regulator of IL-1 beta activity. IL-1ra functions as a competitive antagonist by preventing IL-1 alpha and IL-1 beta from interacting with IL-1 RI. The human IL-1 beta cDNA encodes a 269 aa precursor. A 116 aa propeptide is cleaved intracellularly by the cysteine protease IL-1 beta-converting enzyme (Caspase-1/ICE) to generate the active cytokine. The 17 kDa mature human IL-1 beta b shares 96% aa sequence identity with rhesus and 67-78% with canine, cotton rat, equine, feline, mouse, porcine, and rat IL-1 beta.
Keywords	catabolin;IL1 beta;IL-1 beta;IL-1;IL1B;IL-1b;IL1-BETA;IL-1F2;IL1F2IL-1 beta;interleukin 1; beta;interleukin-1 beta;preinterleukin 1 beta;pro-interleukin-1-beta

GENE INFORMATION

Entrez Gene ID	3553
UniProt ID	P01584