



Rat Anti-Mouse PD-1 (CD279) Monoclonal antibody, clone 29F.1A12 (CABT-L4510)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview

The 29F.1A12 monoclonal antibody reacts with mouse PD-1 (programmed death-1) also known as CD279. PD-1 is a 50-55 kDa cell surface receptor encoded by the *Pdcd1* gene that belongs to the CD28 family of the Ig superfamily. PD-1 is transiently expressed on CD4 and CD8 thymocytes as well as activated T and B lymphocytes and myeloid cells. PD-1 expression declines after successful elimination of antigen. Additionally, *Pdcd1* mRNA is expressed in developing B lymphocytes during the pro-B-cell stage. PD-1's structure includes a ITIM (immunoreceptor tyrosine-based inhibitory motif) suggesting that PD-1 negatively regulates TCR signals. PD-1 signals via binding its two ligands, PD-L1 and PD-L2 both members of the B7 family. Upon ligand binding, PD-1 signaling inhibits T-cell activation, leading to reduced proliferation, cytokine production, and T-cell death. Additionally, PD-1 is known to play key roles in peripheral tolerance and prevention of autoimmune disease in mice as PD-1 knockout animals show dilated cardiomyopathy, splenomegaly, and loss of peripheral tolerance. Induced PD-L1 expression is common in many tumors including squamous cell carcinoma, colon adenocarcinoma, and breast adenocarcinoma. PD-L1 overexpression results in increased resistance of tumor cells to CD8 T cell mediated lysis. In mouse models of melanoma, tumor growth can be transiently arrested via treatment with antibodies which block the interaction between PD-L1 and its receptor PD-1. For these reasons anti-PD-1 mediated immunotherapies are currently being explored as cancer treatments. Like the RMP1-14 and J43 antibodies the 29F.1A12 antibody has been shown to block the binding of PD-1 to its ligands in vivo.

Target	Mouse PD-1 (CD279)
Immunogen	Recombinant PD-1-Ig fusion protein
Isotype	IgG2a, κ
Source/Host	Rat
Species Reactivity	Mouse

Clone	29F.1A12
Purification	Protein G purified. Purity>95%. Determined by SDS-PAGE
Conjugate	Functional Grade
Applications	in vivo blocking of PD-1/PD-L signaling, in vitro PD-1 neutralization, IHC-F, IF, WB, FC
Molecular Weight	150 kDa
Format	0.2 µM filtered liquid. Purified from tissue culture supernatant in an animal free facility
Concentration	Lot specific
Size	5 mg
Buffer	PBS, pH 7.0. Contains no stabilizers or preservatives. [low endotoxin azide-free] Endotoxin level: <2EU/mg (<0.002EU/µg). Determined by LAL gel clotting assay Related dilution buffer: CABT-LB04
Preservative	None
Storage	The antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.
Ship	Wet ice

BACKGROUND

Introduction	This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul 2008]
Keywords	PDCD1;programmed cell death 1;PD1;PD-1;CD279;SLEB2;hPD-1;hPD-I;hSLE1;programmed cell death protein 1;protein PD-1;systemic lupus erythematosus susceptibility 2;

GENE INFORMATION

Official Symbol	programmed cell death 1
------------------------	-------------------------

Synonyms	PDCD1; programmed cell death 1; PD1; PD-1; CD279; SLEB2; hPD-1; hPD-I; hSLE1; programmed cell death protein 1; protein PD-1; systemic lupus erythematosus susceptibility 2;
References	Wang, W., et al. (2018). "RIP1 Kinase Drives Macrophage-Mediated Adaptive Immune Tolerance in Pancreatic Cancer." <i>Cancer Cell</i> 34(5): 757-774 e757. PubMed; Menke, J., et al. (2007). "Programmed death 1 ligand (PD-L) 1 and PD-L2 limit autoimmune kidney disease: distinct roles." <i>J Immunol</i> 179(11): 7466-7477. PubMed; Barber, D. L., et al. (2006). "Restoring function in exhausted CD8 T cells during chronic viral infection." <i>Nature</i> 439(7077): 682-687. PubMed; Liang, S. C., et al. (2003). "Regulation of PD-1, PD-L1, and PD-L2 expression during normal and autoimmune responses." <i>Eur J Immunol</i> 33(10): 2706-2716. PubMed;