



# Mouse Anti-Human TG2 (Beta Barrel 1 Domain) monoclonal antibody, clone YUH32 (CABT-L6035)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Monoclonal antibody to human TG2 (Beta Barrel 1 Domain)
<b>Specificity</b>	Specificity of this clone was determined in WB with human transglutaminases (TG1–TG7, FXIII), TG2 of different species and human TG2 domains recombinantly produced in E. coli. This clone is specific for the Beta Barrel 1 Domain of human TG2. It does not cross-react with other TG2 from various species (human, guinea pig, rat, mouse and dog). It does not cross-react with other human transglutaminases. It does not cross-react with other domains of human TG2.
<b>Immunogen</b>	Human tissue transglutaminase (full length protein with N-terminal hexahistidin-tag) recombinantly produced in insect cells
<b>Isotype</b>	IgG2a
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	YUH32
<b>Purification</b>	The IgG fraction was purified by ion exchange chromatography.
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	WB, IF. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.

<b>Epitope</b>	This clone recognizes the epitope KYLLNLNL
<b>Format</b>	Liquid, Purified
<b>Concentration</b>	Lot specific
<b>Size</b>	200 µg
<b>Buffer</b>	75 mM NaCl, 5 mM Tris, pH7.5, 0.025% sodium azide, 50% glycerol.
<b>Preservative</b>	0.025% sodium azide
<b>Storage</b>	Store at -80°C. If storage at -80°C is not possible, storage at $\leq$ -20°C is recommended. Stable for short term at +4°C.
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	Tissue transglutaminase is a, Ca2+-dependent enzyme (78 kDa) composed by 4 domains: Beta Sheet Domain (fibronectin binding, ~17 kDa), catalytic Core Domain (Cys-His-Asp catalytic triad, Calcium-binding, GTP/GDP-binding, ~37 kDa), Beta Barrel 1 Domain (GTP/GDP-binding, ~14 kDa) and Beta Barrel 2 Domain (~12 kDa). The inactive GTP-bound enzyme is present in a closed conformation, which upon activation by Ca2+ and substrate binding opens like a pocket knife resulting in a longitudinal open conformation
<b>Keywords</b>	Tissue transglutaminase; TG1; TG2; TG3; TG4; TG5; TG6; TG7; keratinocyte transglutaminase; tissue transglutaminase; epidermal transglutaminase; prostate transglutaminase; neuronal transglutaminase