



Rabbit Anti-Human HDAC6 monoclonal antibody, clone KK10-10 (CABT-L845)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Target	HDAC6
Immunogen	Recombinant protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Clone	KK10-10
Purification	Protein A purified.
Conjugate	Unconjugated
Applications	WB, ICC/IF, IHC, IP
Molecular Weight	131 kDa
Cellular Localization	Nucleus, Cytoplasm, Perikaryon, Cell projection.
Positive Control	293, Jurkat, HeLa, HepG2, human kidney tissue.
Format	Liquid
Size	100 µl
Buffer	1×TBS (pH7.4), 1% BSA, 40% Glycerol.
Preservative	0.05% Sodium Azide

Storage

Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

BACKGROUND

Introduction

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, PCAF (p300/CBP associated factor), p300/CBP, HAT1, and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3-6, have been identified as histone deacetylases.

Keywords

CPBHM;FLJ16239;HD 6;HD6;HDAC 6;HDAC6;HDAC6_HUMAN;Histone deacetylase 6 (HD6);Histone deacetylase 6;JM21;JM21;KIAA0901;OTTHUMP00000032398;OTTHUMP00000197663;PPP1R90;Protein phosphatase 1 regulatory subunit 90 antibody

GENE INFORMATION

Entrez Gene ID

[10013](#)

UniProt ID

[A0A024QZ26](#)
