



Rabbit Anti-Cynomolgus TNFRSF1B Polyclonal Antibody (CABT-NS1741)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	Cynomolgus TNFR2/CD120b/TNFRSF1B
Target	TNFRSF1B
Immunogen	Recombinant Cynomolgus TNFR2/CD120b/TNFRSF1B protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Cynomolgus
Conjugate	Unconjugated
Applications	<p>ELISA</p> <p>Recommended dilution:</p> <p>ELISA: 0.1-0.2 µg/mL.</p> <p>This antibody can be used at 0.1-0.2 µg/mL with the appropriate secondary reagents to detect Cynomolgus TNFR2/CD120b/TNFRSF1B.</p> <p>The detection limit for Cynomolgus TNFR2/CD120b/TNFRSF1B is 0.00975 ng/well.</p> <p>Each laboratory should determine an optimum working titer for use in its particular application.</p> <p>Other applications have not been tested but use in such assays should not necessarily be excluded.</p>
Format	Liquid, Purified
Size	50 µl, 100 µl, 200 µl
Buffer	0.2 µm filtered solution in PBS
Preservative	None

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

BACKGROUND

Introduction

Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B), also known as Tumor necrosis factor receptor 2 (TNFR2) or CD120b antigen, is a member of the tumor necrosis factor receptor superfamily. TNFR2/CD120b/TNFRSF1B is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two anti-apoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways. TNFR2/CD120b/TNFRSF1B is not a major contributing factor to the genetic risk of type 2 diabetes, its associated peripheral neuropathy and hypertension and related metabolic traits in North Indians. Tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B) has been reported to be associated with SLE risk in Japanese populations. TNFR2/CD120b/TNFRSF1B serves as a receptor with high affinity for TNFSF2 and approximately 5-fold lower affinity for homotrimeric TNFSF1. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.

Keywords

TNFRSF1B; tumor necrosis factor receptor superfamily, member 1B; p75; TBP11; TNFR; TNFR2; CD120b; TNFR1B; TNFR80; TNF-R75