



Rabbit anti-Arabidopsis thaliana MYC2 (C-term) Polyclonal Antibody (CABT-Z100R)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Immunogen	Antibodies were produced by immunizing animals with a GST-fusion protein containing the C-terminal region of arabidopsis thaliana MYC2.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Arabidopsis thaliana
Purification	Antigen affinity purification
Conjugate	Unconjugated
Applications	WB Recommended dilution: WB: 1:500-1:2,000 (detect endogenous protein*)
Molecular Weight	68 kDa
Preparation	Rabbit polyclonal antibodies were produced by immunizing animals with a GST-fusion protein containing the C-terminal region of arabidopsis thaliana MYC2 (AT1G32640).
Format	Liquid
Concentration	Lot specific
Size	100 µl
Buffer	Supplied in 1 x PBS (pH 7.4), 100 ug/ml BSA, 40% Glycerol, 0.01% NaN ₃ .
Preservative	0.01% NaN ₃

Storage	Store at -20°C. Stable for 6 months from date of receipt.
---------	---

Ship	Wet ice
------	---------

BACKGROUND

Introduction

MYC2 is a common transcription factor of light, abscisic acid (ABA), and jasmonic acid (JA) signaling pathways. Together with MYC3 and MYC4, MYC2 controls subsets of JA-dependent responses. MYC2 is also involved in the regulation of ABA-inducible genes under drought stress conditions in cooperation with MYB2. MYC2 can form complexes with all known glucosinolate-related MYBs to regulate glucosinolate biosynthesis. MYC2 binds to the MYC recognition site (5'-CACATG-3'), the G-box (5'-CACNTG-3') and Z-box (5'-ATACGTGT-3') of the promoters. MYC2 binds directly to the promoters of the transcription factors PLETHORA1 (PLT1) and PLT2 and represses their expression. MYC2 is a negative regulator of blue light-mediated photomorphogenic growth and blue- and far-red-light regulated gene expression. In contrast, MYC2 is a positive regulator of lateral root formation. MYC2 regulates sesquiterpene biosynthesis. MYC2 is subjected to proteasome-dependent proteolysis. The presence of the destruction element (DE) of MYC2 involved in turnover is required for the proper function of MYC2 to regulate gene transcription.

Keywords

Transcription factor MYC2;AtMYC2;Basic helix-loop-helix protein 6;AtbHLH6;bHLH 6;Protein JASMONATE INSENSITIVE 1;R-homologous Arabidopsis protein 1;RAP-1;Transcription factor EN 38;Z-box binding factor 1 protein;bHLH transcription factor bHLH006;rd22BP1;BHLH6;EN38;JAI1;JIN1;RAP1;RD22BP1;ZBF1;At1g32640;F6N18.4

GENE INFORMATION

Gene Name	MYC2
-----------	------

Entrez Gene ID	840158
----------------	------------------------

UniProt ID	Q39204
------------	------------------------