



# Mouse Anti-Horse CXCL10 monoclonal antibody, clone 44-2 (CABT-ZB096)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

Target	CXCL10
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Horse
Clone	44-2
Conjugate	Unconjugated
Applications	FACS, evaluating innate immune responses in horses
Buffer	PBS
Preservative	Sodium azide
Storage	Long time storage is recommended at -20°C.
Ship	Wet ice

## BACKGROUND

Introduction	C-X-C motif chemokine ligand 10 (CXCL10) also known as Interferon gamma-induced protein 10 (IP-10) or small-inducible cytokine B10 is an 8.7 kDa protein that in humans is encoded by the CXCL10 gene. C-X-C motif chemokine 10 is a small cytokine belonging to the CXC chemokine family.
Keywords	CXCL10; chemokine (C-X-C motif) ligand 10; C7; IFI10; INP10; IP-10; crg-2; mob-1; SCYB10;

gIP-10; C-X-C motif chemokine 10; gamma IP10; gamma-IP10; small-inducible cytokine B10; interferon-inducible cytokine IP-10; 10 kDa interferon gamma-induced protein; protein 10 from interferon (gamma)-induced cell line; small inducible cytokine subfamily B (Cys-X-Cys); member 10; NI-0801; HuAIP12

---

## GENE INFORMATION

### Synonyms

CXCL10; chemokine (C-X-C motif) ligand 10; C7; IFI10; INP10; IP-10; crg-2; mob-1; SCYB10; gIP-10; C-X-C motif chemokine 10; gamma IP10; gamma-IP10; small-inducible cytokine B10; interferon-inducible cytokine IP-10; 10 kDa interferon gamma-induced protein; protein 10 from interferon (gamma)-induced cell line; small inducible cytokine subfamily B (Cys-X-Cys); member 10; NI-0801; HuAIP12

---