



Mouse Anti-Human ICOS ligand monoclonal antibody, clone NN18 (CABT-ZB1086)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human ICOS ligand It has no cross-reactivity in ELISA with Human cell lysate (293 cell line).
Target	ICOS
Immunogen	Recombinant Human ICOS Ligand/B7-H2/ICOSLG protein
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	NN18
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA, ELISA(det) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB781 - CABT-ZB1086 This antibody will detect ICOS ligand in antibody pair set. [ABPR-ZB361]
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human ICOS Ligand / B7-H2 / ICOSLG. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	Purified, Liquid

Concentration	Lot specific
Size	50 µL, 100 µL, 200 µL, 1 mL
Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction	Inducible co-stimulator ligand (ICOSL), also known as B7-H2, is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. It is a transmembrane glycoprotein with extracellular IgV and IgC domains and binds to ICOS on activated T cells, thus delivers a positive costimulatory signal for optimal T cell function. The structural features of ICOSL are crucial for its costimulatory function. The present study shows that ICOSL displays a marked oligomerization potential, resembling more like B7-1 than B7-2. B7-H2-dependent signaling may play an active role in a proliferative response rather than in cytokine and chemokine production. The CD28/B7 and ICOS/B7-H2 pathways are both critical for costimulating T cell immune responses. Deficiency in either pathway results in defective T cell activation, cytokine production, and germinal center formation.
---------------------	--

Keywords	IFNA5; interferon, alpha 5; interferon alpha-5; IFN alphaG
-----------------	--

GENE INFORMATION

Synonyms	IFNA5; interferon, alpha 5; interferon alpha-5; IFN alphaG; IeIF G; IFN-alpha-5; interferon alpha-G; interferon alpha-61; INFA5; IFN-alphaG
-----------------	---

Entrez Gene ID	3442
-----------------------	----------------------

UniProt ID	P01569
-------------------	------------------------