



Mouse Anti-Human CEACAM1 monoclonal antibody, clone NN13 (CABT-ZB1103)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human CEACAM1 It has no cross-reactivity in ELISA with Human CEACAM5, Human CEACAM6.
Target	CEACAM1
Immunogen	Recombinant Human CD66a protein
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Clone	NN13
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA, ELISA(det), IHC-P, FC We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB812 - CABT-ZB1103 This antibody will detect CEACAM1 in antibody pair set. [ABPR-ZB393]
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human CEACAM1 / CD66a. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	Purified, Liquid

Concentration	Lot specific
Size	50 µL, 100 µL, 1 mL
Buffer	PBS
Preservative	None
Storage	<p>This antibody can be stored at 2°C-8°C for one month without detectable loss of activity.</p> <p>Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.</p>
Ship	Wet ice

BACKGROUND

Introduction	The carcinoembryonic-antigen-related cell-adhesion molecule (CEACAM) family of proteins has been implicated in various intercellular-adhesion and intracellular-signalling-mediated effects that govern the growth and differentiation of normal and cancerous cells. CEACAM1, also known as biliary glycoprotein I (BGP I) and CD66a, is a member of the carcinoembryonic antigen (CEA) gene family which belongs to the immunoglobulin superfamily. The highly glycosylated CEACAM1 contains one N-terminal V-type Ig-like domain and three C2-type Ig-like domains within its ECD, and one ITIM motif and a calmodulin binding site in the cytoplasmic region. CEACAM1 is a surface glycoprotein expressed on various blood cells, epithelial cells, and vascular cells. It was described as an adhesion molecule mediating cell adhesion via both homophilic and heterophilic manners, and was detected on leukocytes, epithelia, and endothelia. Studies have revealed that CEACAM1 performs actions in multiple cellular processes including tissue differentiation, angiogenesis, apoptosis, metastasis, as well as the modulation of innate and adaptive immune responses.
Keywords	CEACAM1; carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein); BGP; carcinoembryonic antigen-related cell adhesion molecule 1

GENE INFORMATION

Synonyms	CEACAM1; carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein); BGP; carcinoembryonic antigen-related cell adhesion molecule 1; BGP1; CD66a; antigen CD66; CD66a antigen; BGPI
Entrez Gene ID	634
UniProt ID	P13688