



# Mouse Anti-Human NBL1 monoclonal antibody, clone NN17 (CABT-ZB504)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	It reacts with Human NBL1
<b>Target</b>	NBL1
<b>Immunogen</b>	Recombinant Human NBL1/DAND1 Protein
<b>Isotype</b>	IgG
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	NN17
<b>Purification</b>	Protein A purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB504 - CABT-ZB876 This antibody will detect NBL1 in antibody pair set. [ABPR-ZB080]
<b>Preparation</b>	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human NBL1 / DAND1. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
<b>Format</b>	Purified, Liquid
<b>Concentration</b>	Lot specific

<b>Size</b>	50 µL, 100 µL, 200 µL, 1 mL
<b>Buffer</b>	PBS
<b>Preservative</b>	None
<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	The Dan (Differential screening-selected gene aberrative in neuroblastoma, also known as N03) gene was first identified as the putative rat tumor suppressor gene and encodes a protein structurally related to Cerberus and Gremlin in the vertebrates. It is a founding member of the DAN family of secreted proteins, acts as an inhibitor of cell cycle progression, and is closely involved in retinoic acid-induced neuroblastoma differentiation. There are at least five mammalian protein members in the evolutionarily conserved Dan family including DAN, Gremlin/DRM, Cer1 (Cerberus-related), Dante, and PRDC (protein related to DAN and Cerberus), and share the C-terminal cystine-knot motif. As a secreted glycoprotein, DAN is a member of a class of glycoproteins shown to be secreted inhibitors of the transforming growth factor-beta (TGF-beta) and bone morphogenic protein pathways. It binds to BMPs and preventing their interactions with signaling receptor complexes, and accordingly regulates the processes of embryonic development and tissue differentiation. DAN gene product may have an important role in the regulation of the entry of cells into the S phase. Besides, the DAN gene product possesses an ability to revert phenotypes of transformed rat fibroblasts and represents a candidate tumor suppressor gene for neuroblastoma.
<b>Keywords</b>	NBL1; neuroblastoma, suppression of tumorigenicity 1; neuroblastoma suppressor of tumorigenicity 1; D1S1733E

## GENE INFORMATION

<b>Synonyms</b>	NBL1; neuroblastoma, suppression of tumorigenicity 1; neuroblastoma suppressor of tumorigenicity 1; D1S1733E; DAN; DAND1; differential screening selected gene aberrant in neuroblastoma; NB; neuroblastoma candidate region; suppression of tumorigenicity 1
<b>Entrez Gene ID</b>	<a href="#">4681</a>
<b>UniProt ID</b>	<a href="#">P41271</a>