



Rabbit Anti-Human CD21 monoclonal antibody, clone S120 (CABT-ZB576)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human CD21
Target	CR2
Immunogen	Recombinant Human CD21 Protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Clone	S120
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB576 - CABT-ZB930 This antibody will detect CD21 in antibody pair set. [ABPR-ZB153]
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Human CD21.
Format	Purified, Liquid
Concentration	Lot specific
Size	50 µL, 100 µL, 1 mL

Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction	The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD21, also known as Complement component (3d/Epstein Barr virus) receptor 2 and CR2, is a member of the CD system and is a protein involved in complement system. CD21 is present on all mature B-cells and some T-cells and follicular dendritic cells. CD21 on mature B-cells form a complex called the B cell receptor complex with two other membrane proteins, CD19 and CD81. CD21 has a function in the complement system through serving as the cellular receptor specific for ligands such as C3 and C4 which can be attached to foreign macromolecules in order to remove or uptake them. This results in B-cells having enhanced response to the antigen.
---------------------	--

Keywords	CR2; complement component (3d/Epstein Barr virus) receptor 2; CR; C3DR
-----------------	--

GENE INFORMATION

Synonyms	CR2; complement component (3d/Epstein Barr virus) receptor 2; CR; C3DR; CD21; CVID7; SLEB9; complement receptor type 2; EBV receptor; complement C3d receptor
-----------------	---

Entrez Gene ID	1380
-----------------------	----------------------

UniProt ID	P20023
-------------------	------------------------