



# Mouse Anti-Human CCL20 monoclonal antibody, clone NN32 (CABT-ZB597)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	It reacts with Human CCL20
<b>Target</b>	CCL20
<b>Immunogen</b>	Recombinant Human CCL20/MIP-3 alpha/MIP3A Protein
<b>Isotype</b>	IgG
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	NN32
<b>Purification</b>	Protein A purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB597 - CABT-ZB948 This antibody will detect CCL20 in antibody pair set. [ABPR-ZB175]
<b>Preparation</b>	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human CCL20/MIP-3 alpha/MIP3A. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
<b>Format</b>	Purified, Liquid
<b>Concentration</b>	Lot specific

<b>Size</b>	50 $\mu$ L, 100 $\mu$ L, 200 $\mu$ L, 1 mL
<b>Buffer</b>	PBS
<b>Preservative</b>	None
<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	Chemokine (C-C motif) ligand 2 (CCL2) or liver activation regulated chemokine (LARC) or Macrophage Inflammatory Protein-3 (MIP3A) is a small cytokine belonging to the CC chemokine family that attracts immature dendritic cells and memory T lymphocytes, both expressing CCR6. Depending on the cell type, this chemokine was found to be inducible by cytokines (IL-1beta) and by bacterial, viral, or plant products (including LPS, dsRNA, and PMA). MIP3A/CCL2 is Expressed predominantly in the liver, lymph nodes, appendix, peripheral blood lymphocytes, and fetal lung. Low levels of MIP3A/CCL2 has been seen in thymus, prostate, testis, small intestine and colon. As a chemotactic factor, MIP3A/CCL2 attracts lymphocytes and, slightly, neutrophils, but not monocytes. This chemokine may Inhibit proliferation of myeloid progenitors in colony formation assays and it may be involved in formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells. Its C-terminal processed forms have been shown to be equally chemotactically active for leukocytes. Chemokine CCL2 was shown to play a role in colorectal cancer (CRC) pathogenesis.
<b>Keywords</b>	CCL20; chemokine (C-C motif) ligand 20; SCYA20, small inducible cytokine subfamily A (Cys Cys), member 20; C-C motif chemokine 20

## GENE INFORMATION

<b>Synonyms</b>	CCL20; chemokine (C-C motif) ligand 20; SCYA20, small inducible cytokine subfamily A (Cys Cys), member 20; C-C motif chemokine 20; CKb4; exodus 1; LARC; MIP 3a; ST38; exodus-1
<b>Entrez Gene ID</b>	<a href="#">6364</a>
<b>UniProt ID</b>	<a href="#">P78556</a>