



# Rabbit Anti-Mouse SIGNR1 monoclonal antibody, clone S234 (CABT-ZB655)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	It reacts with Mouse SIGNR1
<b>Target</b>	CD209B
<b>Immunogen</b>	Recombinant Mouse SIGNR1/CD209b Protein
<b>Isotype</b>	IgG
<b>Source/Host</b>	Rabbit
<b>Species Reactivity</b>	Mouse
<b>Clone</b>	S234
<b>Purification</b>	Protein A purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB655 - CABT-ZB995 This antibody will detect SIGNR1 in antibody pair set. [ABPR-ZB234]
<b>Preparation</b>	This antibody was obtained from a rabbit immunized with purified, recombinant Mouse SIGNR1 / CD209b.
<b>Format</b>	Purified, Liquid
<b>Concentration</b>	Lot specific
<b>Size</b>	50 µL, 100 µL, 1 mL

<b>Buffer</b>	PBS
<b>Preservative</b>	None
<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD209b, also known as SIGNR1, is a C-type lectin receptor. CD209b is present on most regions of mouse brain and found on microglia and dendritic cells but not on neurons or astrocytes. CD209b is implicated in the recently described SIGNR1 complement activation pathway, which operates against capsular polysaccharides in splenic marginal macrophages. CD209b in rat is homologue to SIGNR1 in mouse, both of which are found to mediate the uptake of dextran or CPS14 within the splenic marginal zone.
<b>Keywords</b>	CD209B; CD209b antigen; CD209 antigen-like protein B; CD209 antigen like protein B

## GENE INFORMATION

<b>Synonyms</b>	CD209B; CD209b antigen; CD209 antigen-like protein B; CD209 antigen like protein B; CD209b antigen; DC SIGN related protein 1; DC SIGNR1; OtB7; DC-SIGNR1; DC-SIGN-related protein 1; OtB7; SIGNR1; mSIGNR1; 1810030I22Rik
<b>Entrez Gene ID</b>	<a href="#">69165</a>
<b>UniProt ID</b>	<a href="#">Q8CJ91</a>