



Mouse Anti-Human TIMP-1 monoclonal antibody, clone NN14 (CABT-ZB662)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human TIMP-1
Target	TIMP1
Immunogen	Human cell-derived TIMP1 protein
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Clone	NN14
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA, ELISA(cap) This antibody will detect TIMP-1 in antibody pair set. [ABPR-ZB241]
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human TIMP1. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	Purified, Liquid
Concentration	Lot specific
Size	50 µL, 100 µL, 1 mL

Buffer	PBS
Preservative	None
Storage	<p>This antibody can be stored at 2°C-8°C for one month without detectable loss of activity.</p> <p>Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.</p>
Ship	Wet ice

BACKGROUND

Introduction	<p>TIMP metallopeptidase inhibitor 1, also known as TIMP-1/TIMP1, Collagenase inhibitor 16C8 fibroblast Erythroid-potentiating activity, TPA-S1TPA-induced proteinTissue inhibitor of metalloproteinases 1, is a natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. TIMP-1/TIMP1 is found in fetal and adult tissues. Highest levels are found in bone, lung, ovary and uterus. Complexes with metalloproteinases and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-1/TIMP1 mediates erythropoiesis in vitro; but, unlike IL-3, it is species-specific, stimulating the growth and differentiation of only human and murine erythroid progenitors. In addition to its inhibitory role against most of the known MMPs, the protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this protein encoding gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This encoding gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction. Complexes with metalloproteinases and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-1/TIMP1 is Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-11, MMP-12, MMP-13 and MMP-16.</p>
Keywords	TIMP1; TIMP metallopeptidase inhibitor 1; EPA; EPO

GENE INFORMATION

Synonyms	TIMP1; TIMP metallopeptidase inhibitor 1; EPA; EPO; HCl; CLGI; TIMP; metalloproteinase inhibitor 1; TIMP-1; collagenase inhibitor; erythroid potentiating activity; erythroid-potentiating activity; fibroblast collagenase inhibitor; tissue inhibitor of metalloproteinases 1; anti-TIMP-1
Entrez Gene ID	7076
UniProt ID	P01033