



Rabbit Anti-Mouse L-selectin monoclonal antibody, clone S134 (CABT-ZB677)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Mouse L-selectin
Target	SELL
Immunogen	Recombinant Mouse CD62L/L-Selectin/SELL Protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Mouse
Clone	S134
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA(cap) This antibody will detect L-selectin in antibody pair set. [ABPR-ZB256]
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Mouse CD62L / L-Selectin / SELL.
Format	Purified, Liquid
Concentration	Lot specific
Size	50 µL, 100 µL, 1 mL
Buffer	PBS

Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction	L-selectin (SELL), also known as CD62L, is a key adhesion molecule that regulates both the migration of leukocytes at sites of inflammation and the recirculation of lymphocytes between blood and lymphoid tissues. It belongs to the selectin family of proteins, and consisting of a large, highly glycosylated, extracellular domain, a single spanning transmembrane domain and a small cytoplasmic tail. L-selectin is the only selectin expressed on leukocytes and mediates a number of leukocyte-endothelial interactions. L-selectin acts as a "homing receptor" for leukocytes to enter secondary lymphoid tissues via high endothelial venules. Ligands present on endothelial cells will bind to leukocyte expressing L-selectin, slowing leukocyte trafficking through the blood, and facilitating entry into a secondary lymphoid organ at that point. L-selectin-mediated lymphocyte recirculation is required for maintaining the appropriate tissue distribution of lymphocyte subpopulations including naïve and effector subsets such as regulatory T cells. In addition, L-selectin-mediated entry into peripheral lymph nodes is required for optimal induction of lymphocyte homeostatic proliferation during lymphopenia. Importantly, L-selectin has been shown to have both adhesive and signaling functions during leukocyte migration. L-selectin has also been shown to mediate leukocyte recruitment during chronic inflammatory and autoimmune diseases and thus is a potential therapeutic target for drug development.
--------------	---

Keywords	SELL; selectin L; TQ1; LAM1
----------	-----------------------------

GENE INFORMATION

Synonyms	SELL; selectin L; TQ1; LAM1; LEU8; LNHR; LSEL; CD62L; LYAM1; PLNHR; LECAM1
Entrez Gene ID	20343
UniProt ID	P18337