



Mouse Anti-Human PDGFRB monoclonal antibody, clone NN17 (CABT-ZB714)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human PDGFRB
Target	PDGFRB
Immunogen	Recombinant Human PDGFRB/CD140b Protein
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	NN17
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB714 - CABT-ZB1039 This antibody will detect PDGFRB in antibody pair set. [ABPR-ZB294]
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human PDGFRB / CD140b. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	Purified, Liquid
Concentration	Lot specific

Size	50 μ L, 100 μ L, 200 μ L, 1 mL
Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction	The cluster of differentiation (CD) system is commonly used as cell markers in Immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD140b, also known as PDGFRB, is a member of the CD system. CD140b is a cell surface tyrosine kinase receptor essential for development interacting with the platelet-derived growth factors (PDGFs) which serves as mitogens for mesenchymal cells. CD140b can bind with platelet-derived growth factor (PDGF)-B, that are secreted by tumors and phosphorylation of PDGFR- β was correlated with depth of cancer invasion at statistically significant level.
---------------------	--

Keywords	PDGFRB; platelet-derived growth factor receptor, beta polypeptide; IMF1; IBGC4
-----------------	--

GENE INFORMATION

Synonyms	PDGFRB; platelet-derived growth factor receptor, beta polypeptide; IMF1; IBGC4; JTK12; PDGFR; CD140B; PDGFR1; PDGFR-1; platelet-derived growth factor receptor beta; PDGFR-beta; PDGF-R-beta; CD140 antigen-like family member B; platelet-derived growth factor receptor 1; beta-type platelet-derived growth factor receptor
-----------------	--

Entrez Gene ID	5159
-----------------------	----------------------

UniProt ID	Q06124
-------------------	------------------------