



# Mouse Anti-Human LDLR/LDL Receptor monoclonal antibody, clone NN13 (CABT-ZB741)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	It reacts with Human LDLR/LDL Receptor
<b>Target</b>	LDLR
<b>Immunogen</b>	Recombinant Human LDLR/LDL R/LDL Receptor Protein
<b>Isotype</b>	IgG1
<b>Source/Host</b>	Mouse
<b>Species Reactivity</b>	Human
<b>Clone</b>	NN13
<b>Purification</b>	Protein A purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB741 - CABT-ZB1059 This antibody will detect LDLR/LDL Receptor in antibody pair set. [ABPR-ZB321]
<b>Preparation</b>	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human LDLR / LDL R / LDL Receptor. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
<b>Format</b>	Purified, Liquid

<b>Concentration</b>	Lot specific
<b>Size</b>	50 $\mu$ L, 100 $\mu$ L, 200 $\mu$ L, 1 mL
<b>Buffer</b>	PBS
<b>Preservative</b>	None
<b>Storage</b>	<p>This antibody can be stored at 2°C-8°C for one month without detectable loss of activity.</p> <p>Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.</p>
<b>Ship</b>	Wet ice

## BACKGROUND

<b>Introduction</b>	LDL Receptor, also known as LDLR, is a mosaic protein that belongs to the Low-density lipoprotein receptor gene family. The low-density lipoprotein receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. LDL Receptor consists of 840 amino acids (after removal of signal peptide) and mediates the endocytosis of cholesterol-rich LDL. Low-density lipoprotein (LDL) is normally bound at the cell membrane and taken into the cell ending up in lysosomes where the protein is degraded and the cholesterol is made available for repression of microsomal enzyme 3-hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase, the rate-limiting step in cholesterol synthesis. At the same time, a reciprocal stimulation of cholesterol ester synthesis takes place. LDL Receptor is a cell-surface receptor that recognizes the apoprotein B100 which is embedded in the phospholipid outer layer of LDL particles. The receptor also recognizes the apoE protein found in chylomicron remnants and VLDL remnants.
---------------------	---

<b>Keywords</b>	LDLR; low density lipoprotein receptor; FH; FHC
-----------------	---

## GENE INFORMATION

<b>Synonyms</b>	LDLR; low density lipoprotein receptor; FH; FHC; LDLCQ2; low-density lipoprotein receptor; LDL receptor; low-density lipoprotein receptor class A domain-containing protein 3
<b>Entrez Gene ID</b>	<a href="#">3949</a>
<b>UniProt ID</b>	<a href="#">P01130</a>