



# Rabbit Anti-Human CEACAM1 monoclonal antibody, clone S114 (CABT-ZB812)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Specificity</b>	It reacts with Human CEACAM1
<b>Target</b>	CEACAM1
<b>Immunogen</b>	Recombinant Human CEACAM-1/CD66a Protein
<b>Isotype</b>	IgG1
<b>Source/Host</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Clone</b>	S114
<b>Purification</b>	Protein A purified
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	ELISA(cap) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB812 - CABT-ZB1103 This antibody will detect CEACAM1 in antibody pair set. [ABPR-ZB393]
<b>Preparation</b>	This antibody was obtained from a rabbit immunized with purified, recombinant Human CEACAM-1 / CD66a.
<b>Format</b>	Purified, Liquid
<b>Concentration</b>	Lot specific
<b>Size</b>	50 µL, 100 µL, 1 mL

Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

## BACKGROUND

**Introduction** The carcinoembryonic-antigen-related cell-adhesion molecule (CEACAM) family of proteins has been implicated in various intercellular-adhesion and intracellular-signalling-mediated effects that govern the growth and differentiation of normal and cancerous cells. CEACAM1, also known as biliary glycoprotein I (BGP I) and CD66a, is a member of the carcinoembryonic antigen (CEA) gene family which belongs to the immunoglobulin superfamily. The highly glycosylated CEACAM1 contains one N-terminal V-type Ig-like domain and three C2-type Ig-like domains within its ECD, and one ITIM motif and a calmodulin binding site in the cytoplasmic region. CEACAM1 is a surface glycoprotein expressed on various blood cells, epithelial cells, and vascular cells. It was described as an adhesion molecule mediating cell adhesion via both homophilic and heterophilic manners, and was detected on leukocytes, epithelia, and endothelia. Studies have revealed that CEACAM1 performs actions in multiple cellular processes including tissue differentiation, angiogenesis, apoptosis, metastasis, as well as the modulation of innate and adaptive immune responses.

**Keywords** CEACAM1; carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein); BGP; carcinoembryonic antigen-related cell adhesion molecule 1

## GENE INFORMATION

**Synonyms** CEACAM1; carcinoembryonic antigen-related cell adhesion molecule 1 (biliary glycoprotein); BGP; carcinoembryonic antigen-related cell adhesion molecule 1; BGP1; CD66a; antigen CD66; CD66a antigen; BGPI

**Entrez Gene ID** [634](#)

**UniProt ID** [P13688](#)