



Mouse Anti-Human DPP2 monoclonal antibody, clone NN14 (CABT-ZB840)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human DPP2
Target	DPP7
Immunogen	Human cell-derived DPPII protein
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	NN14
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA, ELISA(det) We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB442 - CABT-ZB840 This antibody will detect DPP2 in antibody pair set. [ABPR-ZB016]
Preparation	This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human DPPII / DPP7. The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.
Format	Purified, Liquid
Concentration	Lot specific

Size	50 μ L, 100 μ L, 200 μ L, 1 mL
Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction DPP7 (dipeptidylpeptidase 7), also known as DPPII and DPP2, is a post-proline cleaving aminopeptidase expressed in quiescent lymphocytes. Dipeptidyl peptidases (DPPs) have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. It localizes to lysosomes. DPP7 localizes to lysosomes and exists as a homodimer via its leucine zipper motif and is involved in the degradation of oligopeptides. In response to calcium release, it can be secreted in its active form. It is essential for lymphocyte survival, as the inhibition of DPP7 results in quiescent cell apoptosis.

Keywords DPP7; dipeptidyl-peptidase 7; QPP; DPP2

GENE INFORMATION

Synonyms DPP7; dipeptidyl-peptidase 7; QPP; DPP2; DPPII; dipeptidyl peptidase 2; DPP II; carboxytripeptidase; dipeptidyl peptidase II; dipeptidyl-peptidase II; dipeptidyl arylamidase II; dipeptidyl aminopeptidase II; quiescent cell proline dipeptidase

Entrez Gene ID [29952](#)

UniProt ID [Q969X4](#)