



Rabbit Anti-Human CD146/MCAM monoclonal antibody, clone S155 (CABT-ZB943)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human CD146/MCAM It has no cross-reactivity in ELISA with Human CD66a/CEACAM1, Human CD166/alcama, Human CD106/VCAM1, Human CD226/DNAM-1, Human BCAM/CD239.
Target	MCAM
Immunogen	Recombinant Human CD146 protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Clone	S155
Purification	Protein A purified
Conjugate	Unconjugated
Applications	ELISA, ELISA(det), IHC-P, FC, ICC/IF We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB591 - CABT-ZB943 This antibody will detect CD146/MCAM in antibody pair set. [ABPR-ZB168]
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Human CD146 / MCAM extracellular domain.
Format	Purified, Liquid
Concentration	Lot specific

Size	50 µL, 100 µL, 1 mL
Buffer	PBS
Preservative	None
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Ship	Wet ice

BACKGROUND

Introduction	The CD146 antigen, also known as melanoma cell adhesion molecule (MCAM) and MUC18, is an integral membrane glycoprotein belonging to the immunoglobulin superfamily. CD146 contains the characteristic immunoglobulin-like domains (V-V-C2-C2-C2), a transmembrane region, and a short cytoplasmic tail. The CD146 expression is detected in endothelial cells in vascular tissue throughout the body and plays a role in cell adhesion, as well as in cohesion of the endothelial monolayer at intercellular junctions in vascular tissue. As a Ca2+-independent cell adhesion molecule involved in heterophilic cell to cell interactions and a surface receptor, CD146 triggers tyrosine phosphorylation of FYN and PTK2 and subsequently induced signal transduction, proteolysis, or immune recognition. This protein is also expressed predominantly on metastatic lesions and advanced primary tumors, and thus has been suggested to play an important role in tumor progression and the development of metastasis in certain human carcinomas.
---------------------	---

Keywords	MCAM; melanoma cell adhesion molecule; CD146; MUC18
-----------------	---

GENE INFORMATION

Synonyms	MCAM; melanoma cell adhesion molecule; CD146; MUC18; cell surface glycoprotein MUC18; Gicerin; melanoma adhesion molecule; cell surface glycoprotein P1H12; melanoma-associated antigen A32; melanoma-associated antigen MUC18; S-endo 1 endothelial-associated antigen; ABX-MA1
-----------------	--

Entrez Gene ID	4162
-----------------------	----------------------

UniProt ID	P43121
-------------------	------------------------