



Rabbit Anti-Human C1 inhibitor monoclonal antibody, clone S129 (CABT-ZB969)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	It reacts with Human C1 inhibitor It has no cross-reactivity in ELISA with Human SerpinA1/A1AT, Human SerpinA3/SERPINA3/AACT, Human SerpinA5/PROCI, Human SerpinC1 /SERPINC1/ Antithrombin-III, Human SerpinD1/SERPIND1/HC2, Human SerpinF1/SERPINF1/PEDF, Human SerpinF2/SERPINF2, Human SerpinI1/SERPINI1/Protease inhibitor 12.
Target	SERPING1
Immunogen	Recombinant Human SerpinG1 protein
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Human
Clone	S129
Purification	Protein A purified
Conjugate	Unconjugated
Applications	WB, ELISA, ELISA(det), IP We recommend the following for sandwich ELISA (Capture - Detection): CABT-ZB621 - CABT-ZB969 This antibody will detect C1 inhibitor in antibody pair set. [ABPR-ZB200]
Preparation	This antibody was obtained from a rabbit immunized with purified, recombinant Human SerpinG1.
Format	Purified, Liquid

Concentration	Lot specific
Size	50 μ L, 100 μ L, 1 mL
Buffer	PBS
Preservative	None
Storage	<p>This antibody can be stored at 2°C-8°C for one month without detectable loss of activity.</p> <p>Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.</p>
Ship	Wet ice

BACKGROUND

Introduction	Plasma protease C1 inhibitor, also known as C1-inhibiting factor, C1-INH, C1 esterase inhibitor, SERPING1 and C1IN, is a serine proteinase inhibitor (serpin) that regulates activation of both the complement and contact systems. By its C-terminal part (serpin domain), characterized by three beta-sheets and an exposed mobile reactive loop, C1-INH binds, and blocks the activity of its target proteases. The N-terminal end (nonserpin domain) confers to C1-INH the capacity to bind lipopolysaccharides and E-selectin. Owing to this moiety, C1-INH intervenes in regulation of the inflammatory reaction. The heterozygous deficiency of C1-INH results in hereditary angioedema (HAE). Owing to its ability to modulate the contact and complement systems and the convincing safety profile, plasma-derived C1 inhibitor is an attractive therapeutic protein to treat inflammatory diseases other than HAE. Deficiency of C1 inhibitor results in hereditary angioedema, which is characterized by recurrent episodes of localized angioedema of the skin, gastrointestinal mucosa or upper respiratory mucosa. C1 inhibitor may prove useful in a variety of other diseases including septic shock, reperfusion injury, hyperacute transplant rejection, traumatic and hemorrhagic shock, and the increased vascular permeability associated with thermal injury, interleukin-2 therapy and cardiopulmonary bypass.
Keywords	SERPING1; serpin peptidase inhibitor, clade G (C1 inhibitor), member 1; C1IN; C1NH

GENE INFORMATION

Synonyms	SERPING1; serpin peptidase inhibitor, clade G (C1 inhibitor), member 1; C1IN; C1NH; HAE1; HAE2; C1INH; plasma protease C1 inhibitor; serpin G1; C1-inhibiting factor; C1 esterase inhibitor; complement component 1 inhibitor; serine/cysteine proteinase inhibitor clade G member 1
Entrez Gene ID	710
UniProt ID	P05155