



# Human BAG2 blocking peptide (CDBP0559)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Blocking/Immunizing peptide for anti-BAG2 antibody
<b>Antigen Description</b>	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.
<b>Species</b>	Human
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	Apuri, BL, ELISA
<b>Format</b>	Lyophilized powder
<b>Size</b>	100 µg
<b>Preservative</b>	None
<b>Storage</b>	Shipped at ambient temperature, store at -20°C.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">BAG2 BCL2-associated athanogene 2 [ Homo sapiens ]</a>
<b>Official Symbol</b>	BAG2
<b>Synonyms</b>	BAG2; BCL2-associated athanogene 2; BAG family molecular chaperone regulator 2; bcl-2-

associated athanogene 2; BAG-family molecular chaperone regulator-2; dJ417I1.2 (BAG-family molecular chaperone regulator 2); BAG-2; dJ417I1.2; KIAA0576; MGC149462;

Entrez Gene ID	<a href="#">9532</a>
mRNA Refseq	<a href="#">NM_004282</a>
Protein Refseq	<a href="#">NP_004273</a>
UniProt ID	O95816
Chromosome Location	6p12.3-p11.2
Pathway	Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem;
Function	chaperone binding;