



## APAF1 blocking peptide (CDBP5072)

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

<b>Antigen Description</b>	This gene encodes a cytoplasmic protein that initiates apoptosis. This protein contains several copies of the WD-40 domain, a caspase recruitment domain (CARD), and an ATPase domain (NB-ARC). Upon binding cytochrome c and dATP, this protein forms an oligomeric apoptosome. The apoptosome binds and cleaves caspase 9 preproprotein, releasing its mature, activated form. Activated caspase 9 stimulates the subsequent caspase cascade that commits the cell to apoptosis. Alternative splicing results in several transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	Used as a blocking peptide in immunoblotting applications.
<b>Format</b>	Liquid
<b>Concentration</b>	200 µg/mL
<b>Size</b>	0.05 mg
<b>Preservative</b>	None
<b>Storage</b>	-20°C

### GENE INFORMATION

<b>Gene Name</b>	<a href="#">APAF1 apoptotic peptidase activating factor 1 [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	APAF1
<b>Synonyms</b>	APAF1; apoptotic peptidase activating factor 1; CED4; APAF-1; apoptotic protease-activating factor 1

---

Entrez Gene ID	<a href="#">317</a>
mRNA Refseq	<a href="#">NM_001160</a>
Protein Refseq	<a href="#">NP_001151</a>
UniProt ID	O14727
Pathway	Activation of caspases through apoptosome-mediated cleavage; Alzheimer's disease; Alzheimers Disease; Amyotrophic lateral sclerosis (ALS); Apoptosis; Apoptosis Modulation and Signaling; Apoptosis Modulation by HSP70; Apoptotic factor-mediated response
Function	ADP binding; ATP binding; cysteine-type endopeptidase activator activity involved in apoptotic process; identical protein binding; nucleotide binding; protein binding

---