



BCL2L1 blocking peptide (CDBP5179)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Two alternatively spliced transcript variants, which encode distinct isoforms, have been reported. The longer isoform acts as an apoptotic inhibitor and the shorter form acts as an apoptotic activator. [provided by RefSeq, Jul 2008]
----------------------------	--

Conjugate	Unconjugated
Applications	Used as a blocking peptide in immunoblotting applications.
Format	Liquid
Concentration	200 µg/mL
Size	0.05 mg
Preservative	None
Storage	-20°C

GENE INFORMATION

Gene Name	BCL2L1 BCL2-like 1 [Homo sapiens (human)]
Official Symbol	BCL2L1
Synonyms	BCL2L1; BCL2-like 1; BCLX; BCL2L; BCLXL; BCLXS; Bcl-X; bcl-xL; bcl-xS; PPP1R52; BCL-

XL/S; bcl-2-like protein 1; apoptosis regulator Bcl-X; protein phosphatase 1, regulatory subunit 52

Entrez Gene ID	598
mRNA Refseq	NM_001191
Protein Refseq	NP_001182
UniProt ID	Q07817
Pathway	Amyotrophic lateral sclerosis (ALS); Apoptosis; Apoptosis Modulation and Signaling; BH3-only proteins associate with and inactivate anti-apoptotic BCL-2 members; Chronic myeloid leukemia; EPO signaling pathway; HTLV-I infection; IL-3 Signaling Pathway
Function	BH3 domain binding; identical protein binding; protein binding; protein heterodimerization activity; protein homodimerization activity; protein kinase binding