



CLEC2A blocking peptide (CDBP5312)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| | |
|----------------------------|--|
| Antigen Description | CLEC2A belongs to the CLEC2 family of activation-induced, natural killer gene complex-encoded C-type lectin-like receptors (Spreu et al., 2007 [PubMed 18046548]).[supplied by OMIM, May 2008] |
| Conjugate | Unconjugated |
| Applications | Used as a blocking peptide in immunoblotting applications. |
| Format | Liquid |
| Concentration | 200 µg/mL |
| Size | 0.05 mg |
| Preservative | None |
| Storage | -20°C |

GENE INFORMATION

| | |
|------------------------|--|
| Gene Name | CLEC2A C-type lectin domain family 2, member A [Homo sapiens (human)] |
| Official Symbol | CLEC2A |
| Synonyms | CLEC2A; C-type lectin domain family 2, member A; KACL; PILAR; UNQ5792; INPE5792; C-type lectin domain family 2 member A; keratinocyte-associated C-type lectin; proliferation-induced lymphocyte-associated receptor |
| Entrez Gene ID | 387836 |
| mRNA Refseq | NM_001130711 |

| | |
|-----------------------|--|
| Protein Refseq | NP_001124183 |
| UniProt ID | Q6UVW9 |
| Function | carbohydrate binding; protein binding; protein homodimerization activity |