



CXCL12 blocking peptide (CDBP5350)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	This gene encodes a stromal cell-derived alpha chemokine member of the intercrine family. The encoded protein functions as the ligand for the G-protein coupled receptor, chemokine (C-X-C motif) receptor 4, and plays a role in many diverse cellular functions, including embryogenesis, immune surveillance, inflammation response, tissue homeostasis, and tumor growth and metastasis. Mutations in this gene are associated with resistance to human immunodeficiency virus type 1 infections. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2013]
Conjugate	Unconjugated
Applications	Used as a blocking peptide in immunoblotting applications.
Format	Liquid
Concentration	200 µg/mL
Size	0.05 mg
Preservative	None
Storage	-20°C

GENE INFORMATION

Gene Name	CXCL12 chemokine (C-X-C motif) ligand 12 [Homo sapiens (human)]
Official Symbol	CXCL12
Synonyms	CXCL12; chemokine (C-X-C motif) ligand 12; IRH; PBSF; SDF1; TLSF; TPAR1; SCYB12; stromal cell-derived factor 1; intercrine reduced in hepatomas; pre-B cell growth-stimulating factor

Entrez Gene ID	6387
mRNA Refseq	NM_000609
Protein Refseq	NP_000600
UniProt ID	P48061
Pathway	Axon guidance; CXCR4-mediated signaling events; Chemokine receptors bind chemokines; Chemokine signaling pathway; Class A/1 (Rhodopsin-like receptors); Cytokine-cytokine receptor interaction; G alpha (i) signalling events; GPCR downstream signaling
Function	CXCR chemokine receptor binding; chemokine activity; chemokine receptor binding; growth factor activity; receptor binding
