



## DCAF12 blocking peptide (CDBP6242)

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

<b>Antigen Description</b>	This gene encodes a WD repeat-containing protein that interacts with the COP9 signalosome, a macromolecular complex that interacts with cullin-RING E3 ligases and regulates their activity by hydrolyzing cullin-Nedd8 conjugates. [provided by RefSeq, Jul 2009]
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	Used as a blocking peptide in immunoblotting applications.
<b>Format</b>	Liquid
<b>Concentration</b>	200 µg/mL
<b>Size</b>	0.05 mg
<b>Preservative</b>	None
<b>Storage</b>	-20°C

### GENE INFORMATION

<b>Gene Name</b>	<a href="#">DCAF12 DDB1 and CUL4 associated factor 12 [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	DCAF12
<b>Synonyms</b>	DCAF12; DDB1 and CUL4 associated factor 12; CT102; TCC52; WDR40A; KIAA1892; DDB1-and CUL4-associated factor 12; WD repeat domain 40A; cancer/testis antigen 102; WD repeat-containing protein 40A; centrosome-related protein TCC52; testis cancer centrosome-related protein
<b>Entrez Gene ID</b>	<a href="#">25853</a>
<b>mRNA Refseq</b>	<a href="#">NM_015397</a>

**Protein Refseq** [NP\\_056212](#)

---

**UniProt ID** Q5T6F0

---