



WDR92 blocking peptide (CDBP6427)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

| | |
|----------------------------|---|
| Antigen Description | This gene encodes a protein with two WD40 repeat domains thought to be involved in an apoptosis via activation of caspase-3. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2012] |
| Conjugate | Unconjugated |
| Applications | Used as a blocking peptide in immunoblotting applications. |
| Format | Liquid |
| Concentration | 200 µg/mL |
| Size | 0.05 mg |
| Preservative | None |
| Storage | -20°C |

GENE INFORMATION

| | |
|------------------------|--|
| Gene Name | WDR92 WD repeat domain 92 [Homo sapiens (human)] |
| Official Symbol | WDR92 |
| Synonyms | WDR92; WD repeat domain 92; WD repeat-containing protein 92; monad; WD repeat-containing protein Monad |
| Entrez Gene ID | 116143 |
| mRNA Refseq | NM_001256476 |
| Protein Refseq | NP_001243405 |

| | |
|-------------------|--|
| UniProt ID | Q96MX6 |
| Pathway | Activation of BAD and translocation to mitochondria; Activation of BH3-only proteins; Apoptosis; Ca2+ pathway; DARPP-32 events; FCERI mediated Ca+2 mobilization; Fc epsilon receptor (FCERI) signaling; Immune System |
| Function | methylated histone binding; protein binding |