



# Pseudotyped VSV-SARS-CoV-2 S-ΔG-Luciferase (UK variant B.1.1.7, 20I/501Y.V1, Alpha) (COV-PSV19)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Product Overview</b>	Pseudotyped VSV-SARS-CoV-2 S-ΔG-Luciferase (UK variant B.1.1.7, 20I/501Y.V1, Alpha) encodes the antigenomic-sense (or positive-sense) RNA of a replicon-restricted recombinant vesicular stomatitis virus (rVSV) in which the glycoprotein (G) gene has been replaced with the firefly luciferase (Fluc) reporter gene and SARS-CoV-2 spike protein (ΔH69/V70, ΔY144, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H) is incorporated as the membrane protein on the surface of the VSV pseudotyped virus. Because the infectivity of Pseudotyped VSV-SARS-CoV-2 S-ΔG-Luciferase is restricted to a single round of replication, the pseudotypes can be handled using BSL-2 containment practices. The pseudotype VSV particles encode luciferase together with the VSV nucleocapsid (N), phosphoprotein (P), glycoprotein (G), and large polymerase subunit (L) in their pVSV-ΔG vector. When the VSV pseudovirus infects the target cells, luciferase expression is proportional to the number of cells that were infected
<b>Applications</b>	We recommend to use 2-5 uL pseudotyped virus per 1E+04 293T cells for in vitro assay. Due to differences in cell status, the best infection conditions and MOI should be determined by the end user. The virus can be diluted with cell culture medium if needed.
<b>Size</b>	1 ml
<b>Storage</b>	Store at -80°C. Multiple freeze/thaw cycles not recommended. When using the virus, transfer the virus from the -80 ° C refrigerator and melt it in an ice bath.
<b>Ship</b>	Frozen on dry ice