



## Anti-TDO2 (full length) polyclonal antibody (CPBT-47955MH)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

Product Overview	Mouse Polyclonal antibody to Human TDO2.
Antigen Description	Tryptophan 2,3-dioxygenase (EC 1.13.11.11) plays a role in catalyzing the first and rat-limiting step in the kynurenine pathway, the major pathway of tryptophan metabolism.
Immunogen	Full length human TDO2 protein
Isotype	IgG
Source/Host	Mouse
Species Reactivity	Human
Purification	Protein A purified
Conjugate	Unconjugated
Applications	WB, ELISA
Sequence Similarities	Belongs to the tryptophan 2,3-dioxygenase family.
Format	Liquid
Size	50 μg
Buffer	Preservative: NoneConstituents: 1X PBS, pH 7.2
Preservative	None
Storage	Shipped at $4^{\circ}$ C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

## **GENE INFORMATION**

Gene Name	TDO2 tryptophan 2,3-dioxygenase [ Homo sapiens ]
Official Symbol	TDO2
Synonyms	TDO2; tryptophan 2,3-dioxygenase; TDO; TPH2; 3-dioxygenase; T23O_HUMAN; TDO 2; TDO; tdo2; TO; TPH2; TRPO; Tryptamin 2 3 dioxygenase; Tryptamin 2; Tryptophan 2 3 dioxygenase; Tryptophan 2; Tryptophan oxygenase; Tryptophan pyrrolase; Tryptophanase; TO; tryptophanase; tryptophan oxygenase; tryptophan pyrrolase; tryptamin 2,3-dioxygenase; TRPO;
Entrez Gene ID	<u>6999</u>
Protein Refseq	NP_005642
UniProt ID	<u>P48775</u>
Chromosome Location	4q31-q32
Pathway	Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Monoamine Transport, organism-specific biosystem; NAD biosynthesis II (from tryptophan), conserved biosystem; Tryptophan catabolism, organism-specific biosystem; Tryptophan metabolism, organism-specific biosystem; Tryptophan metabolism, organism-specific biosystem; Tryptophan metabolism, conserved biosystem; tryptophan degradation III (
Function	amino acid binding; heme binding; metal ion binding; oxidoreductase activity; oxygen binding; tryptophan 2,3-dioxygenase activity;