



Human HEXIM1 peptide (DAG-P0044)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Antigen Description	Expression of this gene is induced by hexamethylene-bis-acetamide in vascular smooth muscle cells. This gene has no introns. [provided by RefSeq, Jul 2008]
Specificity	Ubiquitously expressed with higher expression in placenta. HEXIM1 and HEXIM2 are differentially expressed. Expressed in endocrine tissues.
Conjugate	Unconjugated
Sequence Similarities	Belongs to the HEXIM family.
Format	Liquid
Preservative	None
Storage	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. Information available upon request.

GENE INFORMATION

Gene Name	HEXIM1 hexamethylene bis-acetamide inducible 1 [Homo sapiens (human)]
Official Symbol	HEXIM1
Synonyms	HEXIM1; hexamethylene bis-acetamide inducible 1; CLP1; EDG1; HIS1; MAQ1; protein HEXIM1; HMBA-inducible; menage a quatre 1; cardiac lineage protein 1; menage a quatre protein 1; estrogen down-regulated gene 1 protein; hexamethylene bis-acetamide inducible 1; hexamethylene bisacetamide-inducible protein; hexamethylene bis-acetamide-inducible protein 1; hexamethylene-bis-acetamide-inducible transcript 1;
Entrez Gene ID	10614

mRNA Refseq	NM_006460.2
Protein Refseq	NP_006451.1
UniProt ID	O94992
Chromosome Location	17q21.31
Function	cyclin-dependent protein serine/threonine kinase inhibitor activity; protein binding; snRNA binding;