



## IL17C peptide (DAG-P0674)

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

<b>Antigen Description</b>	The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells. [provided by RefSeq, Jul 2008]
<b>Purity</b>	70 - 90% by HPLC.
<b>Conjugate</b>	Unconjugated
<b>Sequence Similarities</b>	Belongs to the IL-17 family.
<b>Format</b>	Liquid
<b>Preservative</b>	None
<b>Storage</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. Information available upon request.

### GENE INFORMATION

<b>Gene Name</b>	<a href="#">IL17C interleukin 17C [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	IL17C
<b>Synonyms</b>	IL17C; interleukin 17C; CX2; IL-17C; interleukin-17C; cytokine CX2;
<b>Entrez Gene ID</b>	<a href="#">27189</a>
<b>mRNA Refseq</b>	<a href="#">NM_013278.3</a>
<b>Protein Refseq</b>	<a href="#">NP_037410.1</a>

<b>UniProt ID</b>	Q9P0M4
<b>Chromosome Location</b>	16q24
<b>Pathway</b>	IL17 signaling pathway, organism-specific biosystem;
<b>Function</b>	cytokine activity;