



# Human PSMA5 peptide (DAG-P1886)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

<b>Antigen Description</b>	The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been found for this gene. [provided by RefSeq, Dec 2010]
----------------------------	--

<b>Specificity</b>	Expressed in fetal brain (at protein level).
<b>Conjugate</b>	Unconjugated
<b>Applications</b>	Neut
<b>Sequence Similarities</b>	Belongs to the peptidase T1A family.
<b>Format</b>	Liquid
<b>Preservative</b>	None
<b>Storage</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">PSMA5 proteasome (prosome, macropain) subunit, alpha type, 5 [ Homo sapiens (human) ]</a>
<b>Official Symbol</b>	PSMA5
<b>Synonyms</b>	PSMA5; proteasome (prosome, macropain) subunit, alpha type, 5; PSC5; ZETA; proteasome

subunit alpha type-5; macropain zeta chain; proteasome zeta chain; macropain subunit zeta; proteasome component 5; proteasome subunit zeta; proteasome alpha 5 subunit; multicatalytic endopeptidase complex zeta chain;

Entrez Gene ID	<a href="#">5686</a>
mRNA Refseq	<a href="#">NM_001199772.1</a>
Protein Refseq	<a href="#">NP_001186701.1</a>
UniProt ID	P28066
Chromosome Location	1p13
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kapp
Function	protein binding; threonine-type endopeptidase activity;