



Recombinant Canine Pancreatic Lipase (DAG-WT1206)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Species	Canine
Purity	> 90% as determined by SDS-PAGE
Conjugate	His
Applications	Immunoassays
Molecular Weight	51.5 kDa
Format	Liquid
Size	1 mg
Buffer	Tris/PBS-based buffer
Preservative	None
Storage	Store at -80°C. Avoid multiple freeze/thaw cycles.
Ship	Dry ice

BACKGROUND

Introduction

Pancreatic lipase, also known as pancreatic triacylglycerol lipase, is secreted from the pancreas, and is the primary lipase (enzyme) that hydrolyzes (breaks down) dietary fat molecules in the human digestive system, converting triglyceride substrates found in ingested oils to monoglycerides and free fatty acids. Triacylglycerol + H2O diacylglycerol + a fatty acid anion Bile salts secreted from the liver and stored in gallbladder are released into the

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

duodenum where they coat and emulsify large fat droplets into smaller droplets, thus increasing the overall surface area of the fat, which allows the lipase to break apart the fat more effectively. The resulting monomers (2 free fatty acids and one 2-monoacylglycerol) are then moved by way of peristalsis along the small intestine to be absorbed into the lymphatic system by a specialized vessel called a lacteal. This protein belongs to pancreatic lipase family.

Keywords

Pancreatic lipase; PNLIP; Canine

GENE INFORMATION

UniProt ID

A0A8I3QDI3