



Mouse Monocyte Chemotactic Protein 1 (DAG285)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Recombinant Mouse Monocyte Chemotactic Protein-1 (MCP-1) (CCL2/JE). Recombinant Mouse MCP-1 is a single, non-glycosylated polypeptide chain containing 125 amino acids and having a molecular weight of 13,803 Da. The sequence of the first five N-terminal amino acids is: D-Asp-Asp-Asp-Asp-Asp.
Antigen Description	Chemokine(C-C motif) ligand 2 (CCL2) is a small cytokine belonging to the CC chemokine family that is also known as monocyte chemotactic protein-1 (MCP-1) and small inducible cytokine A2. CCL2 recruits monocytes, memory T cells, and dendritic cells to sites of tissue injury, infection, and inflammation.
Species	Mouse
Conjugate	Unconjugated
Applications	The activity is calculated by the ability to chemoattract Balb/c mouse spleen MNCs at 1.0-20 ng/ml. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays.
Format	Purified, Lyophilized. Reconstitute using sterile deionized water to a concentration 100 µg/ml. Further dilutions can be made in other aqueous buffers.
Concentration	Lot specific (OD280nm) (prior to lyophilization)
Buffer	Not applicable.
Preservative	None
Storage	2-8°C short term, -20°C long term

BACKGROUND

Introduction	This gene is one of several cytokine genes clustered on the q-arm of chromosome 17.
--------------	---

Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. [provided by RefSeq, Jul 2013]

Keywords	CCL2; chemokine (C-C motif) ligand 2; HC11; MCAF; MCP1; MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30; C-C motif chemokine 2; small-inducible cytokine A2; monocyte secretory protein JE; monocyte chemotactic protein 1; monocyte chemoattractant protein 1; monocyte
----------	--
