



## Anti-IL1RL1 monoclonal antibody, clone 08304 (DCABY-3957)

This product is for research use only and is not intended for diagnostic use.

## PRODUCT INFORMATION

Antigen Description	The Interleukin 1 receptor family comprises at least nine members. All family members possess three immunoglobulin (Ig)-like domains in their extracellular region. Most members also have an intracellular TIR (Toll-like receptor/IL-1 receptor) signaling domain that is also conserved in the Toll-like receptor family. A subtype of the IL-1 R family, SIGIRR (single Ig domain-containing IL-1 R-related molecule) differs from the other nine members by having only one Ig domain. ST2L, also known as IL-1 R4, is a transmembrane glycoprotein in the interleukin-1 receptor family. ST2L is expressed on mast cells and Th2 cells as well as on stressed cardiac myocytes and airway epithelia. ST2L associates with IL-1R AcP to form a receptor complex for IL-33. Alternative splicing generates ST2, a secreted isoform that functions as an inhibitor of IL-33 signaling. Within the ECD, mouse ST2L shares 68% aa sequence identity with human ST2L.
Specificity	Detects human ST2/IL-1R4 in ELISAs andWestern blots. In Western blots,no cross-reactivity with recombinant mouseST2L, recombinant human (rh)IL-1RI, rhIL-1 RII, rhIL-1 RA, rhIL-1 alpha, or rhIL-1 beta is observed.
Immunogen	S. frugiperda insect ovarian cell line Sf 21-derived recombinant human ST2/IL-1 R4. Lys19-Phe328 Accession Number BAA02233
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	08304
Purification	Protein A or G purified from hybridoma culture supernatant
Conjugate	Unconjugated

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

Applications	Western Blot, ELISA Capture (Matched Pair), Neutralization
Format	Liquid
Size	500 μg
Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.
Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  12 months from date of receipt, -20 to -70 °C as supplied.  1 month, 2 to 8 °C under sterile conditions after reconstitution.  6 months, -20 to -70 °C under sterile conditions after reconstitution.

## **GENE INFORMATION**

Gene Name	IL1RL1 interleukin 1 receptor-like 1 [ Homo sapiens (human) ]
Official Symbol	IL1RL1
Synonyms	IL1RL1; interleukin 1 receptor-like 1; T1; ST2; DER4; ST2L; ST2V; FIT-1; IL33R; interleukin-1 receptor-like 1; growth stimulation-expressed; interleukin 1 receptor-related protein; homolog of mouse growth stimulation-expressed;
Entrez Gene ID	9173
Protein Refseq	NP 001269337
UniProt ID	Q01638
Chromosome Location	2q12
Function	cytokine receptor activity; interleukin-1 receptor activity; interleukin-33 binding; interleukin-33 receptor activity; protein binding; receptor signaling protein activity;