



## Anti-CXCL11 monoclonal antibody, clone 98439 (DCABY-3984)

This product is for research use only and is not intended for diagnostic use.

## **PRODUCT INFORMATION**

Antigen Description	Interferon-inducible T cell a chemoattractant (I-TAC), also known as SCYB9B, H174 and beta-R1, is a non-ELR motif-containing CXC chemokine. I-TAC shares 36% and 37% amino acid sequence homology with IP-10 and MIG, respectively. I-TAC is expressed at low levels in normal tissues, including thymus, spleen and pancreas.
Specificity	Detects human CXCL11/I-TAC in ELISAs and Western blots. In ELISAs, this antibody does not cross-react with recombinant human (rh) CXCL5, 7, 8, 9, or 10.In Western blots, this antibody shows less than 1% cross-reactivity with rhCXCL1, 2, 3, 5, 6, 7, IL-8/CXCL8, 9, 10, 13, CXCL12/SDF-1 alpha, CXCL12/SDF-1 beta, rmCXCL1, 2, 6, 9, 10, CXCL12/SDF-1 alpha, rmCXCL13, rrCXCL1, CXCL3/CINC-2 alpha, CXCL3/CINC-2 beta, or rpIL-8/CXCL8.
Immunogen	E. coli-derived recombinant human CXCL11/I-TAC. Phe22-Phe94 Accession Number O14625
Isotype	lgG2a
Source/Host	Mouse
Species Reactivity	Human
Clone	98439
Purification	Protein A or G purified from ascites
Conjugate	Unconjugated
Applications	Western Blot, ELISA Capture (Matched Pair), Neutralization
Format	Liquid
Size	500 μg

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Email: info@creative-diagnostics.com

© Creative Diagnostics All Rights Reserved

Buffer	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.
Preservative	None
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	12 months from date of receipt, -20 to -70 °C as supplied.
	1 month from date of receipt, 2 to 8 °C, reconstituted.
	6 months from date of receipt, -20 to -70 °C, reconstituted.

## **GENE INFORMATION**

Gene Name	CXCL11 chemokine (C-X-C motif) ligand 11 [ Homo sapiens (human) ]
Official Symbol	CXCL11
Synonyms	CXCL11; chemokine (C-X-C motif) ligand 11; IP9; H174; IP-9; b-R1; I-TAC; SCYB11; SCYB9B; C-X-C motif chemokine 11; beta-R1; small inducible cytokine B11; small-inducible cytokine B11; interferon gamma-inducible protein 9; interferon-inducible T-cell alpha
Entrez Gene ID	<u>6373</u>
Protein Refseq	NP 005400
UniProt ID	<u>014625</u>
Chromosome Location	4q21.2
Pathway	CXCR3-mediated signaling events; Chemokine receptors bind chemokines; Chemokine signaling pathway; Class A/1 (Rhodopsin-like receptors); Cytokine-cytokine receptor interaction; G alpha (i) signalling events; GPCR downstream signaling; GPCR ligand binding;
Function	CXCR3 chemokine receptor binding; chemokine activity; heparin binding; protein binding;